

LOUISIANA PLANT PATHOLOGY

DISEASE IDENTIFICATION AND MANAGEMENT SERIES

LSU
AgCenter
Research & Extension

Grain Spotting and Pecky Rice

Damage by many fungi including *Cochliobolus miyabeanus* (Ito and Kuribayashi) Drechs. Ex Dastur, *Curvularia* spp., *Fusarium* spp., *Microdochium oryzae* (Hashioka and Yokogi) Samuels and I.C. Halett, *Sarocladium oryzae* (Sawada) W. Gams and D. Hawksworth and other fungi

Many fungi infect developing grain and cause spots and discoloration on the hulls or kernels. Damage by the rice stink bug, *Oebalus pugnax* F., also causes discoloration of the kernel (Figure 1).

Kernels discolored by fungal infections or insect damage commonly are called pecky rice or peck. With the disorder, the endosperm of the grain shows what can range from small areas to being almost totally rotten (Figure 2).

This complex disorder involves many fungi (Figure 3), the white-tip nematode and insect damage. High winds at the early heading stage may cause similar symptoms. Heading applications of stink bug insecticides and fungicides targeted at other more important diseases will reduce peck.



Figure 2. Pecky rice



Figure 1. Rice stink bug feeding



Figure 3. Fungi growing from pecky rice

Visit our website: www.lsuagcenter.com

Authors

Don Groth, Ph.D., Professor
Rice Research Station

Clayton Hollier, Ph.D., Professor
Department of Plant Pathology and
Crop Physiology

Louisiana State University Agricultural Center, William B. Richardson, Chancellor
Louisiana Agricultural Experiment Station, David J. Boethel, Vice Chancellor and Director
Louisiana Cooperative Extension Service, Paul D. Coreil, Vice Chancellor and Director

Pub. 3112 (online only) 12/10
The LSU AgCenter is a statewide campus of the LSU System and provides equal opportunities in programs and employment.