

SOIL TEST INFORMATION SHEET NO. V-340

Prepared by Extension Horticulturist James Boudreaux and modified by Research Coordinator Christine Coker

Greens (Mustard, Turnips, Spinach, Parsley, Kale and Collards)

1. Greens are sensitive to strongly acid soils. Soils with a pH below 6.0 should be limed with agricultural limestone well ahead of planting. If magnesium is low, apply dolomitic limestone. Apply lime 2-6 months before planting. Spinach is more sensitive to low pH soils than other greens. Spinach should be grown on soils with a pH of 6.0-6.8.
2. Apply about half the nitrogen (50 lbs./acre) and all the phosphorus and potassium prior to planting.
3. Sidedress greens with 25 to 30 pounds nitrogen per acre when greens are 4 to 6 inches tall, 3 to 4 weeks after planting. Use higher rates of fertilizer on double drill greens.
4. Boron deficiency is common in turnip roots and all greens when grown on soils with a pH below 5.5. Boron deficiency in turnip is expressed as brown rings inside the root. Boron deficiency can be prevented by applying 3 pounds of actual boron per acre with preplant fertilizer. Apply 10-25 pounds of borax per acre or 5-12 pounds of Solubor per acre with preplant fertilizer. Foliar applications of borax (2-5 pounds per 100 gals.) or Solubur (1 to 1 1/2 lbs./100 gals.) can be used. Always apply foliar treatments in the early morning hours to the underside of young rapidly growing plants for best results. Generally 2 or 3 applications 7 to 10 days apart are sufficient to avoid boron deficiency.
5. Greens grown on fields following other vegetable crops need only a nitrogen sidedressing.
6. Contact your county agent for additional information and help in your fertilization program. The agent also receives a copy of this report for the parish office files.